



Rehabilitation Protocol for MPFL Reconstruction

This protocol is intended to guide clinicians through the post-operative course for MPFL reconstruction. This protocol is time based (dependent on tissue healing) as well as criterion based. Specific intervention should be based on the needs of the individual and should consider exam findings and clinical decision making. The timeframes for expected outcomes contained within this guideline may vary based on surgeon's preference, additional procedures performed, and/or complications. If a clinician requires assistance in the progression of a post-operative patient, they should consult with the referring surgeon.

The interventions included within this protocol are not intended to be an inclusive list. Therapeutic interventions should be included and modified based on the progress of the patient and under the discretion of the clinician.

Considerations with concomitant procedures:

Many different factors influence the post-operative MPFL reconstruction rehabilitation outcomes, including additional procedure such as tibial tuberosity osteotomy (TTO). It is recommended that clinicians collaborate closely with the referring physician regarding early range of motion, weight bearing status, and use of assistive devices.

Post-operative considerations:

If you develop a fever, excessive drainage from incision, severe heat and/or redness along incision, uncontrolled pain, or any other symptoms that concern you please call your doctor.

PHASE I: IMMEDIATE POST-OP (0-2 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> Protect surgical site Reduce swelling, minimize pain Restore full extension, gradually improve flexion ≥ 90 deg Minimize arthrogenic muscle inhibition, re-establish quad control, regain full active extension Patient education <ul style="list-style-type: none"> Keep your knee straight and elevated when sitting or laying down. Do not rest with a towel placed under the knee
Weight Bearing	<p><i>Walking</i></p> <ul style="list-style-type: none"> Initially brace locked, PWB (0-1 week) → WBAT with crutches (per MD recommendation) May start walking without crutches as long as there is no increased pain, effusion, and proper gait When climbing stairs, make sure you are leading with the non-surgical side when going up the stairs, make sure you are leading with the crutches and surgical side when going down the stairs
Interventions	<p><i>Swelling Management</i></p> <ul style="list-style-type: none"> Ice, compression, elevation (check with MD re: cold therapy) Retrograde massage Ankle pumps <p><i>Range of motion/Mobility</i></p> <ul style="list-style-type: none"> PROM Heel slides with towel Low intensity, long duration extension stretches: prone hang, heel prop Seated hamstring/calf stretch <p><i>Strengthening</i></p> <ul style="list-style-type: none"> Calf raises

	<ul style="list-style-type: none"> • Quad sets <ul style="list-style-type: none"> ○ NMES high intensity (2500 Hz, 75 bursts) supine knee extended 10 sec/50 sec, 10 contractions, 2x/wk during sessions—use of clinical stimulator during session, consider home units distributed immediate post op • Straight leg raise <ul style="list-style-type: none"> ○ **Do not perform straight leg raise if you have a knee extension lag • Hip abduction • Standing hamstring curl
Criteria to Progress	<ul style="list-style-type: none"> • Knee extension ROM 0 deg • Quad contraction with superior patella glide and full active extension • Able to perform straight leg raise without lag

PHASE II: INTERMEDIATE POST-OP (3-6 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> • Continue to protect surgical site • Maintain full extension, restore full flexion (contralateral side) • Normalize gait • Patient education
Weight Bearing	<p><i>Walking</i></p> <ul style="list-style-type: none"> • WBAT: May unlock brace when able to perform straight leg raise without lag • Discontinue use of brace after 6 wks (or per surgeon) and when gait is normalized
Additional Interventions <i>*Continue with Phase I interventions</i>	<p><i>Range of motion/Mobility</i></p> <ul style="list-style-type: none"> • Stationary bicycle • Gentle patellar mobilizations: superior/inferior and medial/lateral *Not necessary unless stiffness present <p><i>Strengthening</i></p> <ul style="list-style-type: none"> • Adductor strengthening: hook lying ball squeezes, SLR adduction, bridging with ball squeeze • Ball squats, wall slides, mini squats from 0-60 <p><i>Balance/proprioception</i></p> <ul style="list-style-type: none"> • Single leg standing balance (knee slightly flexed) static progressed to dynamic and level progressed to unsteady surface
Criteria to Progress	<ul style="list-style-type: none"> • No swelling (Modified Stroke Test) • Flexion ROM > 90 deg • Extension ROM equal to contra lateral side

PHASE III: LATE POST-OP (7-12 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> • Continue to protect surgical site • Maintain full ROM • Safely progress strengthening • Promote proper movement patterns • Avoid post exercise pain/swelling • Avoid activities that produce pain at repair site
Weight Bearing	<ul style="list-style-type: none"> • FWB without assistive device
Additional Interventions <i>*Continue with Phase I-II Interventions</i>	<p><i>Range of motion/Mobility</i></p> <ul style="list-style-type: none"> • Gentle stretching all muscle groups: prone quad stretch, standing quad stretch, standing hip flexor stretch <p><i>Cardio</i></p> <ul style="list-style-type: none"> • ~8 weeks: Elliptical, stair climber, flutter kick swimming, pool jogging <p><i>Strengthening</i></p> <ul style="list-style-type: none"> • Gym equipment: leg press machine, seated hamstring curl machine and hamstring curl machine, hip abductor and adductor machine, hip extension machine, roman chair, seated calf machine

	<p>**The following exercises to focus on proper control with emphasis on good proximal stability</p> <ul style="list-style-type: none"> Proximal Strengthening: Double leg bridge, bridge with feet on physio ball, single leg bridge, lateral band walk, standing clamshell/fire hydrant, hamstring walkout, TA brace with UE and LE progression Squat to chair Lateral lunges Romanian deadlift (single and double leg) Single leg progression: single leg press, slide board lunges: retro and lateral, split squats, step ups and step ups with march, lateral step-ups, step downs, single leg squats, single leg wall slides/sit Lateral band walks <p><i>Balance/proprioception</i></p> <ul style="list-style-type: none"> Progress single limb balance including perturbation training
Criteria to Progress	<ul style="list-style-type: none"> No effusion/swelling/pain after exercise Normal gait ROM equal to contra lateral side Quad/HS/glut index $\geq 70\%$; HHD mean or isokinetic testing @ 60d/s

PHASE IV: TRANSITIONAL (13-16 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> Maintain full ROM Safely progress strengthening Promote proper movement patterns Avoid post exercise pain/swelling Avoid activities that produce pain
Additional Interventions <i>*Continue with Phase II-III interventions</i>	<p><i>Strengthening</i></p> <ul style="list-style-type: none"> Progress intensity (weight) and volume (repetitions) of exercises <p><i>Plyometric activities</i></p> <ul style="list-style-type: none"> Bilateral FWB plyometrics progressed to single leg plyometrics <p><i>Balance/proprioception</i></p> <ul style="list-style-type: none"> Progress single limb balance including perturbation training
Criteria to Progress	<ul style="list-style-type: none"> Clearance from MD and ALL milestone criteria below have been met <u>Functional Assessment</u> <ul style="list-style-type: none"> Quad/HS/glut index $\geq 80\%$; HHD mean or isokinetic testing @ 60d/s Hamstring/Quad ratio $\geq 66\%$ Hop Testing $\geq 80\%$ compared to contra lateral side, demonstrating good landing mechanics

PHASE V: EARLY RETURN TO SPORT (3-5 MONTHS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> Safely progress strengthening Safely initiate sport specific training program Promote proper movement patterns Avoid post exercise pain/swelling Avoid activities that produce pain at graft donor site
Additional Interventions <i>*Continue with Phase II-IV interventions</i>	<p><i>Strengthening</i></p> <ul style="list-style-type: none"> Progress intensity (weight) and volume (repetitions) of exercises <p><i>Interval running program</i></p> <ul style="list-style-type: none"> <u>Return to Running Program</u>

	<p><i>Progress to plyometric and agility program (with functional brace if prescribed)</i></p> <ul style="list-style-type: none"> ○ <u>Agility and Plyometric Program</u>
Criteria to Progress	<ul style="list-style-type: none"> • Clearance from MD and ALL milestone criteria below have been met • Completion jog/run program without pain/effusion / swelling • <u>Functional Assessment</u> <ul style="list-style-type: none"> ○ Quad/HS/glut index $\geq 95\%$; HHD mean or isokinetic testing @ 60d/s ○ Hamstring/Quad ratio $\geq 66\%$ ○ Hop Testing $\geq 95\%$ compared to contra lateral side, demonstrating good landing mechanics • <u>Lysholm</u> $>90\%$ • <u>KOOS-sports questionnaire</u> $>90\%$ • <u>International Knee Committee Subjective Knee Evaluation</u> >93 • <u>Psych Readiness to Return to Sport (PRRS)</u> • <u>Kujala</u> > 90

PHASE VI: UNRESTRICTED RETURN TO SPORT (6+ MONTHS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> • Continue strengthening and proprioceptive exercises • Symmetrical performance with sport specific drills • Safely progress to full sport
Additional Interventions <i>*Continue with Phase II-V interventions</i>	<ul style="list-style-type: none"> • Multi-plane sport specific plyometrics program • Multi-plane sport specific agility program • Include hard cutting and pivoting depending on the individuals' goals • Non-contact practice → Full practice → Full play (~6-7 mo)
Criteria to Progress	<ul style="list-style-type: none"> • Last stage, no additional criteria

--	--

References:

Saper MG, Fantozzi P, Bompadre V, et al. Return-to-sport testing after medial patellofemoral ligament reconstruction in adolescent athletes. *Orthop J Sports Med* 2019;7: 2325967119828953.

Manske RC, Prohaska D. Rehabilitation following medial patellofemoral ligament reconstruction for patellar instability. *Int J Sports Phys Ther* 2017;12:494-511.

Cosgarea AJ, Johnson K, McGee TG, et al. Rehabilitation after medial patellofemoral ligament reconstruction. *Sports Med Arthrosc Rev* 2017;25:105-113.

Clark D, Walmsley K, Schranz P, et al. Tibial tuberosity transfer in combination with medial patellofemoral ligament reconstruction: Surgical technique. *Arthrosc Tech* 2017;6:591-597.

Hinckel BB, Gobbi RG, Kaleka CC, et al. Medial patellotibial ligament and medial patellomeniscal ligament: Anatomy, imaging, biomechanics, and clinical review. *Knee Surg Sports Traumatol Arthrosc* 2018;26:685-696.

Ahmad CS, Lightsey HM, Popkin CA, et al. Rehabilitation variability following medial patellofemoral ligament reconstruction. *Phys Sportsmed* 2018;46:441-448